



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$PR$  perpendicular to the opposite side of the angle. Mark off a distance equal to  $2\overline{OP}$  on a straight-edge passing through  $O$ , and adjust the straight-edge so that one mark falls on line  $PQ$  at  $M$ , and the other on line  $PR$  at  $N$ . Draw the line  $NOM$ , which will trisect the angle.

*Proof:* Complete the rectangle  $PMKN$ , and draw the other diagonal  $PK$ . Then  $\overline{OP} = \frac{1}{2}\overline{MN} = \frac{1}{2}\overline{PK} = \overline{PC} = \overline{CM}$ .

Hence,  $\theta = \alpha + \phi = \beta + \phi = \gamma + \delta + \phi = 2\delta + \phi = 2\phi + \phi = 3\phi$ .

## DEPARTMENTS.

### SOLUTIONS OF PROBLEMS.

#### ALGEBRA.

283. Proposed by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Solve  $w+x+y+z=4a$ ,  $w^2+x^2+y^2+z^2=4a^2+4b^2$ ,  $w^3+x^3+y^3+z^3=4a^3+12ab^2$ ,  $w^4+x^4+y^4+z^4=4a^4+4b^4+4c^4+24a^2b^2$ .

Solution by DR. L. E. DICKSON, Associate Professor of Mathematics, The University of Chicago.

The following method applies equally well to the corresponding equations with arbitrary constant terms. We are given  $s_1, s_2, s_3, s_4$ , where  $s_n$  is the sum of the  $n$ th powers of  $w, x, y, z$ . Hence the latter are, by Newton's identities, the roots of the following quartic:

$$\xi^4 - 4a\xi^3 + (6a^2 - 2b^2)\xi^2 + (4ab^2 - 4a^3)\xi + a^4 + b^4 - 2a^2b^2 - c^4 = 0.$$

To obtain the reduced quartic, set  $\xi = \eta + a$ . Then

$$\eta^4 - 2b^2\eta^2 + b^4 - c^4 = 0, \quad (\eta^2 - b^2)^2 = c^4.$$

Hence, the 24 sets of solutions are given by the arrangements of  $a \pm \sqrt{(b^2 \pm c^2)}$ .

Similarly solved by G. B. M. Zerr and J. Scheffer.

#### GEOMETRY.

312. Proposed by F. H. SAFFORD, Ph. D., The University of Pennsylvania, Philadelphia, Pa.

A variable circle passes through a fixed point and is tangent to a given circle. If a diameter of the first circle passes through the fixed point, find the locus of its other extremity.